

## Neisseria Meningitidis

Neisseria meningitidis, often called meningococcus, is a gram negative diplococci that can cause meningitis and life threatening sepsis. Neisseria are fastidious organisms that require nutrient supplementation to grow in laboratory cultures. Specifically, they grow on chocolate agar and can be isolated on Thayer-Martin media, sometimes called VPN agar. Thayer-Martin media is an agar plate containing vancomycin, polymyxin, and nystatin along with other nutrients that facilitate the growth of Neisseria species while inhibiting the growth of other organisms. Characteristically, Neisseria are oxidase positive and maltose-fermenting. *N. meningitidis* can be distinguished from *N. gonorrhoeae* because *N. meningitidis* ferments glucose in addition to maltose while *N. gonorrhoeae* only ferments glucose. These organisms exist as normal flora in the nasopharynx of 5-15% of adults and is spread through the exchange of saliva or other respiratory secretions. Virulence mechanisms include lipooligosaccharide (LOS) within the cell membrane of *N. meningitidis*, which acts as an endotoxin and can cause shock and destruction of red blood cells. Other virulence mechanisms include a polysaccharide capsule that is anti-phagocytic and aids in the evasion of the host immune response. Symptoms can rapidly progress from fatigue to meningitis with fever, headache, and neck stiffness. If toxic levels of the endotoxin reach the bloodstream, meningococcal sepsis can develop causing blood vessels to rupture and vital organs to rapidly shut down. Complications include Waterhouse-Friderichsen syndrome with adrenal insufficiency, organ failure, low blood pressure, disseminated intravascular coagulation with widespread petechial rash, and death. Treatment includes ceftriaxone and rifampin prophylaxis.



PLAY PICMONIC

### Characteristics

#### Gram-Negative

[Graham-cracker Negative-devil](#)

This organism stains negative on Gram stain due to relatively thin peptidoglycan layer in the cell wall.

#### Diplococci

[Double-cockeyes](#)

Diplococci are round-shaped bacteria that typically occur as two joined cells.

#### Chocolate Agar

[Chocolate-bar](#)

Neisseria are fastidious organisms that require nutrient supplementation to grow in laboratory cultures. Specifically, they grow on chocolate agar which is an enriched growth medium containing red blood cells that have been lysed, providing growth factors like NAD and hemin. The agar is named for the chocolate color of the medium and does not actually contain chocolate.

#### Thayer-Martin Media

[Thief driving aston Martin car](#)

Thayer-Martin media is an agar plate containing vancomycin, polymyxin, nystatin and specific nutrients that facilitate the growth of Neisseria species while inhibiting the growth of other organisms.

## VPN

### [VPN on License Plate](#)

Thayer-Martin media is also called VPN agar, standing for vancomycin, polymyxin, and nystatin. These antibiotics along with specific nutrients facilitate the growth of *Neisseria* species while inhibiting the growth of other organisms.

## Oxidase-Positive

### [Ox-daisy](#)

An oxidase test is used to determine if bacteria produce certain cytochrome c oxidases to help differentiate bacteria. *Neisseria* are oxidase-positive bacteria.

## Maltose Fermentation

### [Malt-liquor Fern](#)

*Neisseria meningitidis* can be distinguished from *Neisseria gonorrhoeae* because *N. meningitidis* ferments both glucose and maltose while *N. gonorrhoeae* only ferments glucose.

## Glucose Fermentation

### [Glue-bottle Fern](#)

*Neisseria* can be distinguished from other gram-negative bacteria because they are glucose-fermenting. Both *Neisseria gonorrhoeae* and *Neisseria meningitidis* are glucose fermenting.

## Endotoxin

### [Unpopped-toxic-balloon](#)

Virulence mechanisms include lipooligosaccharide (LOS), also known as endotoxin, within the cell membrane of *N. meningitidis*, which instigates a systemic inflammatory response with shock and destruction of red blood cells.

## Lipooligosaccharide (LOS) in Membrane

### [Lips-sack Capsule in Membrane](#)

LOS is a smaller form of the more common bacterial virulence factor lipopolysaccharide (LPS). LOS is found within the bacterial cell membrane and its function is to maintain membrane stability. LOS is highly antigenic and when released into the bloodstream it causes a systemic inflammatory response.

## Polysaccharide Capsule

### [Polly-sack Capsule](#)

*N. meningitidis* virulence mechanisms include a polysaccharide capsule that is anti-phagocytic and aids in the evasion of the host immune response.

## IgA Protease

### [\(IgA\) Apple-goblin with Propeller-ace](#)

An IgA protease is an enzyme that cleaves certain amino acid sequences of proteins including immunoglobulin A. *Neisseria meningitidis* releases IgA proteases which destroy IgA, leading to increased pathogenicity. Other IgA protease producers include *Streptococcus pneumoniae* and *Haemophilus influenzae* type B.

## Disease

## Waterhouse Friderichsen Syndrome

### [Water-house](#)

This syndrome is defined as adrenal failure due to massive hemorrhage into the adrenal glands caused by overwhelming meningococemia and disseminated intravascular coagulation.

## Adrenal Insufficiency

### Damaged Adrenal gland

Sepsis caused by *Neisseria meningitidis* can lead to massive hemorrhage into one or both adrenal glands, causing rapidly developing adrenocortical insufficiency with loss of cortisol and aldosterone production. This is also called Waterhouse-Friderichsen syndrome.

## Petechial Rash

### Tiki-mask with a Rash

Petechia are small red or purple spots on the body caused by broken capillary blood vessels. A petechial rash is present in approximately 75% of meningococcal sepsis caused by widespread destruction of blood vessels.

## Treatments

### Ceftriaxone

#### Chef-tri-axes

Ceftriaxone is a third generation cephalosporine used in the treatment of *N. meningitidis*.

### Rifampin Prophylaxis

#### Ref-amp Purple-axes

Rifampin is an antibiotic commonly used to treat tuberculosis. People who have had close contact with an individual who has meningococcal infection can be offered rifampin to eradicate the bacteria and prevent disease.